2465 - CPS1 Pharmacv:

CPS1 - Jenny O'Donnell User: Thu, Jan 12, 2012 09:49 Last login:

## Search Protocols Reports Change password Manage profile Help Logout

## Born 11-Mar-1977 (34y) Gender Male CHI No. 777777777 BISCUIT, Bertie Patient Details Last Modified On 12-Jan-2012 By CPS1 Address V Phone and email HRM Methotrexate Process: Concordance > Interactions & precautions > Adverse reactions > Monitoring > Review Adverse reactions: side effects and toxicity Is the patient aware of the common side effects of The most common side effects are: methotrexate? Yes No • Skin (e.g. urticaria, acne, photosensitivity) Is the patient aware of the side effects that occur if they 0 are having a reaction to methotrexate? · Haematopoietic reactions (e.g. anaemia, pneumonia, Yes No septicaemia) Is the patient aware of what to do if they are suffering 0 0 from these signs? · Alimentary tract (e.g. nausea, vomiting, gingivitis, Yes No pharyngitis, intestinal ulceration) Is the patient aware that adverse reactions should be 0 0 · Hepatic (e.g. acute hepatitis or cirrhosis) reported? Yes No • Urogenital (e.g. vaginal discharge or ulcers, cystitis, dysuria) · Pulmonary (e.g. acute pulmonary oedema, fibrosis) CNS (e.g. headache, drowsiness or blurred vision) · Cardiac (hypotension, deep vein thrombosis) · Check the patient's understanding of the side effects of Methotrexate using the information booklet as a prompt. Ask the patient if they are experiencing any side effects. · Refer them to their GP if side effects appear severe. · Record any care issues in the patient's care plan and agree desired outcomes and actions. Signs of methotrexate toxicity or intolerance are: · Breathlessness · Dry persistent cough · Severe vomiting or diarrhoea · Sore throat or mouth ulcers · Unexplained bleeding or bruising · Abdominal discomfort · Dark urine Actions: Check the patient's understanding of the signs of methotrexate toxicity using the information booklet. Remind the patient that they should tell the pharmacist, GP or specialist if they experience any side effects that could indicate toxicity. · Immediately refer the patient to their GP if they are showing signs of toxicity, contacting their GP to alert them. Patients with signs of pneumonitis must stop methotrexate and seek urgent hospital attention.

agree desired outcomes and actions.

Back - Interactions & precautions

via yourself or by them self reporting.

Next - Monitoring

Advise the patient to report any adverse drug reactions through the MHRA Yellow Card Reporting Scheme either

Record any care issues in the patient's care plan and

Save & Review

return to Methotrexate Review Page