### Patient Group Direction for treatment of Herpes Zoster (Shingles) in patients aged 18 years and over

#### **Patient assessment form**

Patient name and address (including	Click or tap here to enter text.	Date of Birth /CHI:	Click or tap here to enter text.		
postcode):		Sex	М		F 🗆
Date of assessment:	Click or tap to enter a date.	Patient is aware that GP will be informed:	YES		NO 🗆

## Patient clinical picture and related appropriate actions

Clinical features/symptom assessment	Yes	No	Actions
Is patient over 18 years of age?			If NO, do not treat with this PGD. Refer to GP/OOH/ED as appropriate.
Does the rash affect a single dermatome?			If NO, do not treat with this PGD. Refer to GP/OOH/ED as appropriate.
Has rash been present for less than 72 hours?			If NO, do not treat with this PGD. Refer to GP/OOH/ED as appropriate.
Is shingles rash affecting areas other than those relating to dermatomes T1 – L2 e.g. extending to around eyes?			
Is patient already taking antiviral medication?			
Known hypersensitivity to aciclovir or any excipients?			
Does the patient have impaired gastrointestinal absorption e.g. Crohn's disease, ulcerative colitis?			
Does patient have acute diarrhoea and vomiting where aciclovir absorption could be impaired?			If YES to any of the exclusion
Is the patient immunocompromised? E.g. auto-immune disease, current chemotherapy or immunosuppressant medication?			criteria , do not treat with this PGD.
Is the patient pregnant?			Refer to GP/OOH/ED as appropriate.
Is patient systemically unwell?			
Known moderate to severe renal impairment? (eGFR <25mL/minute/1.73m <sup>2)</sup> ?			
Is this recurrent shingles? (Immunocompetent patient with a history of 2 or more episodes in last 12 months)			
Concomitant use of interacting medication?			
Has informed consent to treatment been obtained?			If NO, patient is unable to receive treatment.

# Preparation options and supply method

Medicine and strength (Dispersible tablets strictly limited to those unable to swallow standard tablets)	Regimen	Supply method
Aciclovir 800 mg tablets	ONE tablet FIVE times daily spread evenly throughout the day during waking hours (usually at 4 hourly intervals) x 35	PGD via NHS Pharmacy First Scotland
Aciclovir 400 mg tablets	TWO tablets FIVE times daily spread evenly throughout the day during waking hours (usually at 4 hourly intervals) x 70	
Symptomatic management	Appropriate analgesia – paracetamol	NHS Pharmacy First Scotland, OTC or existing supply

### Patient advice checklist

Advice	Provided (tick as appropriate)
How to take medication – with water, regularly and complete the course	
Ensure adequate fluid intake whilst taking aciclovir tablets	
Expected duration of symptoms - to seek medical assistance if symptoms worsen or are not resolving within 7 days	
Patient information leaflet relating to the medication is given to the patient	
Common side effects of medication e.g. nausea, vomiting, diarrhoea and abdominal pain, taste disturbance, photo sensitivity, pruritus, urticaria, fever, tiredness and occasionally headaches or dizziness.	
Check patient has access to symptomatic relief (use of analgesia – paracetamol)	
Avoid sharing of towels and clothes	
Maintain good hand hygiene	
Wear loose fitting clothes to minimise irritation	
Avoid use of topical creams and adhesive dressings as they can cause irritation and delay rash healing	
Person with shingles is infectious until all the vesicles have crusted over (usually 5-7 days after rash onset)	
Avoid contact with others wherever possible, if the rash is weeping and can't be covered. If the lesions have dried or can be covered, this is not necessary	
Person who has not had chicken pox or the varicella vaccine can catch chicken pox from person with shingles (if possible, avoid pregnant women, immunocompromised people and babies younger than 1 month old)	

### Communication

Contact made with	Details (include time and method of communication)
Patient's General Practice (details)	Click or tap here to enter text.

# Details of medication supplied and pharmacist supplying under the PGD

Medication supplied	Click or tap here to enter text.		
Batch number	Click or tap here to enter text.	Expiry date Click or tap to enter a date.	
Print name of pharmacist	Click or tap here to enter text.		
GPhC registration details	Click or tap here to enter text.		
Signature of pharmacist			

#### Patient Group Direction for treatment of Herpes Zoster (Shingles) in patients aged 18 years and over

#### **Notification of supply from community pharmacy**

#### **CONFIDENTIAL WHEN COMPLETED**

Data protection confidentiality note: this message is intended only for the use of the individual or entity to whom it is addressed and may contain information that is privileged, confidential and exempt from disclosure under law. If the reader of this message is not the intended recipient, you are hereby notified that any dissemination, distribution or copying of this communication is strictly prohibited.

GP name	Click or tap here to enter text.	Pharmacy Stamp	/ Address details	
GP practice address	Click or tap here to enter text.			
	Click or tap here to enter text.			
The following patient h	nas attended this pharmacy for			
assessment and potent	tial treatment of Herpes Zoster (Shingles)			
Patient name	Click or tap here to enter text.			
Date of birth/CHI	Click or tap here to enter text.	Pharmacist name		
Patient address	Click or tap here to enter text.	Click or tap here to enter text.		
	Click or tap here to enter text.	GPhC number Click or tap here to enter text.		
Postcode	Click or tap here to enter text.	DateClick or tap to enter a date.		
times daily	iven a 7 day course of aciclovir 800 mg five		]	
•	iven a 7 day course of aciclovir 800 mg five			
	iven self-care advice only		]	
·	le for treatment via PGD for the following			
reasons and has been referred:				
Click or tap here to enter	text.			
Your patient has been a	dvised to contact the practice if symptoms fa	il to resolve followin	g treatment.	
You may wish to include	this information in your patient records.			
Patient consent: I can conf	irm that the information is a true reflection of my indi	ividual circumstances	Consent	
and I give my consent to allow a pharmacist working under the terms of NHS Pharmacy First Scotland			received	
	oriate advice and/or treatment for me. I also give my p			
the pharmacist to pass, to my own GP, details of this consultation and any advice given, or treatment provided. I have been advised that some of the information may be used to assess the uptake of the				
1 · ·	illy anonymous and not be attributable to any individu		_	

This form should now be sent to the patient's GP and a copy retained in the pharmacy