

Please refer to the Summary of Product Characteristics (SPC) and the Co-Amoxiclav PGD in addition to this form.

**ORAL CO-AMOXICLAV FOR TREATMENT OF SKIN INFECTIONS ASSOCIATED WITH INJECTION SITE
COMPLICATIONS E.G. CELLULITIS AND WOUND INFECTIONS
CONSULTATION FORM**

PATIENT NAME.....

DOB/CHI No.....

ADDRESS.....

POSTCODE.....

DATE AND TIME OF CONSULTATION.....

Exclusion Criteria for treatment under the PGD: (if yes to any of the below please refer)

- Patient under 18 years old
- If the area is purulent (immediate referral to medical staff is necessary)
- Hypersensitivity to the active substances, to any of the penicillins or to any of the excipients. (see SPC and PIL for details)
- History of a severe immediate hypersensitivity reaction (e.g. anaphylaxis) to another beta-lactam agent (e.g. a cephalosporin, carbapenem or monobactam)
- History of jaundice/hepatic impairment due to amoxicillin/clavulanic acid
- Severe hepatic disease
- Renal impairment
- Acute sore throat/tonsillitis
- Current treatment with methotrexate, oral typhoid vaccine, probenecid. N.B. This list is not exhaustive. Please check the BNF and refer to a doctor if necessary
- No valid consent to treatment according to NHS Fife policy

Medication counselling and advice

- Take at regular intervals and complete the course unless otherwise directed. ☐
- For best results take at the start of a meal. ☐
- Tablets should be swallowed whole. Do not crush or chew them. ☐
- Seek further medical advice if condition worsens or fails to improve with treatment. ☐

Follow Up/Referral

- Ask the patient to return for review at 7 days, if symptoms recur then refer to Addictions Service doctor, OOH or GP.
- In the best interest to the patient, carry out a review even if they return after 7 days and refer if necessary.
- The patient may be referred to a doctor at any stage, if this is necessary, in the professional opinion of the Pharmacist.
- Patients should be referred to the doctor if treatment proves to be ineffective in relieving the symptoms

Treatment given as per PGD	Co-Amoxiclav 625mg (500/125) tabs, 1 TDS for 7 days Or Co-Amoxiclav 250/62.5mg in 5ml suspension, 10ml TDS for 7 days
Pharmacist Name	
Pharmacist Signature	
Date	
Pharmacy stamp/ Contractor Code	

Retain consultation checklist for your own records only