

Doxycycline 100mg capsules

GG&C PGD ref no: 2026/2904

YOU MUST BE AUTHORISED BY NAME, UNDER THE CURRENT VERSION OF THIS PGD BEFORE YOU ATTEMPT TO WORK ACCORDING TO IT

Change history

Date	Version number	Update
17/11/23	6	<i>Transfer of PGD onto new Template Update to number of supplies by a Community Pharmacist in 3 month period Update to Cautions section Update to Advice to patient/carer section</i>
04/11/25	7	Updated with new reviewer signatories

Doxycycline 100mg capsules

Clinical Condition	
Indication:	Infective exacerbation of Chronic Obstructive Pulmonary Disease (COPD) when patient has allergy to penicillin and/or clinically appropriate in line with local rescue medication guideline/ protocols
Inclusion criteria:	<ul style="list-style-type: none"> • Patients aged 18 and over • Definite diagnosis of COPD • Patient has COPD “self-management plan” agreed with HC professional and presents COPD Rescue Medication card to Community Pharmacy • Patient reports with infective exacerbation characterised by development or increase in sputum purulence and one or more of the following <ul style="list-style-type: none"> ◦ increase in shortness of breath ◦ increase in sputum volume <p>NB – there is no need for a physical assessment to supply, however you may wish to complete physical assessments to calculate a NEWS score if a patient seems poorly and need onward referral if you can.</p>
Exclusion criteria:	<ul style="list-style-type: none"> • Course of doxycycline within the last month with no resolution of symptoms • More than 2 supplies by community pharmacist in any 3-month period as part of COPD rescue meds service or 3 supplies in 12 months from any prescriber/route • Increased breathing rate ≥ 20 breaths/min • Systemic inflammatory response syndrome criteria e.g. temperature greater than $>36^{\circ}\text{C}$, heart rate >90 beats per minute and other symptoms. • Known allergy to doxycycline, tetracyclines or any of the excipients • Long term use of azithromycin • Known bronchiectasis • Patient has a known increased INR • Pregnancy or Breast Feeding • Known immunosuppression including Acute lymphocytic leukaemia, chronic lymphocytic leukaemia, cytomegalovirus infection • Patients with rare hereditary problems of fructose intolerance, glucose galactose malabsorption or sucrose-isomaltose insufficiency

Doxycycline 100mg capsules

	<ul style="list-style-type: none"> • Known myasthenia gravis • Known systemic lupus erythematosus (SLE) • Patients receiving the following medications: methotrexate, oral ciclosporin, oral retinoids, penicillins, Lithium, oral typhoid vaccine • Patients taking ergotamine or methysergide • Patients taking drugs known to increase the metabolism of doxycycline (e.g. phenytoin, barbiturates, primidone, carbamazepine, rifampicin) • Known hepatic impairment • Signs and symptoms of a severe exacerbation (e.g. new/worsening confusion, marked breathlessness/tachypnoea, new onset cyanosis/peripheral oedema, rapid symptom onset)
Cautions/Need for further advice/Circumstances when further advice should be sought from the prescriber:	<p>If the patient is taking warfarin, they should be advised to have their INR checked within 7 days of commencing antibiotic.</p> <p>Absorption of doxycycline may be impaired by concurrently administered sucralfate or antacids containing aluminium, calcium, magnesium or other drugs containing these cations including Quinapril; oral zinc, iron salts or bismuth preparations.</p> <p>The usual advice within NHS GGC is for the patient to withhold the interacting medication for the duration of the course (5 days in this case).</p> <p>NB: The interaction with doxycycline and iron has been seen even with the previous dose separation advice. If the patient is on regular iron it should be stopped for the short duration of the doxycycline course.</p> <p>Use in caution with patients who have alcohol dependence or who are receiving potentially hepatotoxic drugs.(See BNF)</p> <p>Use in caution with patients who have photosensitivity. Avoid exposure to sunlight or sun lamps</p>
Action if patient declines or is excluded:	<p>Refer to prescriber</p>
Referral arrangements for further advice / cautions:	<p>Refer to GP, OOH or NHS24</p>

Doxycycline 100mg capsules

Drug Details	
Name, form & strength of medicine:	Doxycycline 100mg capsules
Route/Method of administration:	Oral
Dosage (include maximum dose if appropriate):	200mg on first day, then 100mg daily thereafter
Frequency:	Once daily
Duration of treatment:	5 days
Maximum or minimum treatment period:	5 days
Quantity to supply/administer:	6
Supply, Administer or Both:	Supply only
▼ Additional Monitoring:*	No
Legal Category:	POM
Is the use outwith the SPC:**	No
Storage requirements:	Store below 25 °C

* The black triangle symbol has now been replaced by European “additional monitoring” (▼)

** Summary of Product Characteristics

Doxycycline 100mg capsules

<p>Warnings including possible adverse reactions and management of these:</p>	<p>Gastrointestinal symptoms are usually mild and seldom necessitate discontinuation of treatment.</p> <p>Headache and/or visual disturbance may indicate benign intracranial hypertension (treatment should be discontinued).</p> <p>Other common side effects include: Photosensitivity reaction, rash including maculopapular and erythematous rashes, Henoch-Schonlein purpura, Urticaria Hypersensitivity (including anaphylactic shock, anaphylactic reaction, anaphylactoid reaction, exacerbation of systemic lupus erythematosus, serum sickness) Peripheral oedema, hypotension, Pericarditis, Tachycardia</p> <p>Patient to see GP, OOH or NHS 24 if side effects problematic.</p> <p>Use the Yellow Card System to report adverse drug reactions. Yellow Cards and guidance on its use are available at the back of the BNF or online at http://yellowcard.mhra.gov.uk/</p>
<p>Advice to patient/carer including written information provided:</p>	<p>Explain treatment and course of action.</p> <p>Advise pt to avoid any OTC meds such as antacids, multivitamins etc without checking if suitable to take with doxycycline.</p> <p>Complete the prescribed course.</p> <p>Swallow the capsules whole with plenty of water in either the resting or standing position and well before going to bed for the night to reduce the likelihood of oesophageal irritation and ulceration.</p> <p>Advise pt to avoid direct exposure to sunlight or sun lamps due to potential photosensitivity reactions.</p> <p>Remind patient to contact GP Practice or anticoagulant service to arrange to have INR checked within 7 days of commencing antibiotic.</p> <p>Visual disturbances such as blurring of vision may occur during treatment with doxycycline and in such cases please advise patients that they must refrain from driving or operating machinery.</p> <p>Give patient a copy of relevant patient information leaflet, if appropriate.</p>

Doxycycline 100mg capsules

	If condition worsens or symptoms persist then seek further medical advice.
Monitoring (if applicable):	N/A
Follow up:	Patients not improving after a few days of starting antibiotic course or if any deterioration should be advised to contact GP or OOH service

Doxycycline 100mg capsules

Staff Characteristics

Professional qualifications:	Those registered health care professionals that are listed and approved in legislation as able to operate under patient group directions and have current registration
Specialist competencies or qualifications:	<p>Has undertaken appropriate training to carry out clinical assessment of patient leading to diagnosis that requires treatment according to the indications listed in this PGD.</p> <p>Has undertaken appropriate training for working under PGDs for the supply and administration of medicines.</p>
Continuing education & training:	The practitioner should be aware of any change to the recommendations for the medicine listed. It is the responsibility of the individual to keep up-to-date with continued professional development.

Referral Arrangements and Audit Trail

Referral arrangements	<i>As per local arrangements/national guidelines, SBAR or equivalent to GP practice within 1 week as per flow chart.</i>
Records/audit trail:	<ul style="list-style-type: none"> • Patient's name, address, date of birth and consent given • Contact details of GP • Diagnosis • Dose, form administered and batch details • Advice given to patient (including side effects) • Signature/name of staff who administered or supplied the medication, and also, if relevant, signature/name of staff who removed/discontinued the treatment • Details of any adverse drug reaction and actions taken including documentation in the patient's medical record • Referral arrangements (including self-care)





References/Resources and comments:

Notes:
 SPC – Summary of Product Characteristics
 BNF – British National Formulary
[GGC Medicines - Antibiotic Allergy and Interactions](#)
 NICE - [Overview | Chronic obstructive pulmonary disease in over 16s: diagnosis and management | Guidance | NICE](#)

Doxycycline 100mg capsules

This Patient Group Direction must be agreed to and signed by all healthcare professionals involved in its use. The original signed copy will be held at Pharmacy Services, Clarkston Court, 56 Busby Road, Glasgow. The PGD must be easily accessible in the clinical setting.

Organisation: NHS Greater Glasgow & Clyde

Professionals drawing up PGD/Authors	
	Designation and Contact Details
<p>*Name: Lorna Brown</p>  <p>Signature: _____ Date: 08/01/2026</p>	<p>Designation: Advanced Pharmacist, CPDT</p> <p>E-mail address: lorna.brown8@nhs.scot</p>
<p>Name: Andrew Christopherson</p>  <p>Signature: _____ Date: 07/01/2026</p>	<p>Designation: Lead Pharmacist Glasgow City HSCP North West</p> <p>E-mail address: Andrew.Christopherson@nhs.scot</p>
<p>Name: Stuart Sutton</p>  <p>Signature: _____ Date: 01/12/2025</p>	<p>Designation: Clinical Director, Renfrewshire HSCP</p> <p>E-mail address: Stuart.Sutton@nhs.scot</p>
<p>Name: Emma McCahill</p>  <p>Signature: _____ Date: 28/01/2026</p>	<p>Designation: Community Pharmacist IP representative</p> <p>E-mail address: emma.mccahill@nhs.scot</p>
<p>Name: _____</p> <p>Signature: _____ Date: _____</p>	<p>Designation: _____</p> <p>E-mail address: _____</p>


* **Lead Author**

** **Antimicrobial Pharmacist if appropriate**


Doxycycline 100mg capsules

AUTHORISATION:


NHSGG&C PGD Sub-Committee of ADTC

Chairman in BLOCK CAPITALS	Signature:	Date:
Dr Craig Harrow		03/12/2025

NHSGG&C PGD Sub-Committee of ADTC


Chief Nurse, North Sector, NHS GG&C in BLOCK CAPITALS	Signature:	Date:
Kevin McAuley		03/12/2025


Pharmacist representative of PGD Sub-Committee of ADTC

Name: in BLOCK CAPITALS	Signature:	Date:
Elaine Paton		03/12/2025


Antimicrobial use

If the PGD relates to an antimicrobial agent, the use must be supported by the NHS GG&C Antimicrobial Management Team (AMT). A member of this team must sign the PGD on behalf of the AMT.

Microbiology approval	Name: Scott Gillen	Designation: Antimicrobial Pharmacist
	 Signature: _____ Date: 03/12/2025 (on behalf of NHS GG&C AMT)	

NHS Greater Glasgow & Clyde Patient Group Direction (PGD) for Community Pharmacists	
Doxycycline 100mg capsules	

Local Authorisation:

Service Area for which PGD is applicable: Community Pharmacy			
Record/Audit Trail	There must be appropriate records kept and maintained by the pharmacist to enable verification of service provision and training requirements, and provide information for internal and external audit and evaluation purposes.		
Nominated individual who agrees to keep list of practitioners operating under the PGD current up to date (Lead Professional):			
Name:	Signature:	Designation:	Date:
Rachel Bruce Email contact address: rachel.bruce.8@nhs.scot		Lead Pharmacist Community Care	29/01/2026

PGDs do not remove inherent professional obligations or accountability

It is the responsibility of each professional to practice only within the bounds of their own competence and in accordance with the General Pharmaceutical Council Standards for Pharmacy Professionals.


Name of Pharmacist		GPhC No	
---------------------------	--	----------------	--

I have read and understood the Patient Group Direction. I acknowledge that it is a legal document and agree to provide these medicines only in accordance with this PGD.

<i>Sign:</i>		<i>Date:</i>
--------------	--	--------------

ONCE YOU HAVE SIGNED THE PGD, YOU MUST COMPLETE THE ELECTRONIC FORM (LINK BELOW)

[PGDs - Greater Glasgow and Clyde \(office.com\)](https://www.nhs.uk/medicines/patient-group-directions/pgds-greater-glasgow-and-clyde-office-com)

NHS Greater Glasgow & Clyde Patient Group Direction (PGD) for Community Pharmacists	
Doxycycline 100mg capsules	

Patient Group Direction Audit Form
Form for the audit of compliance with PGD or PGDs

To ensure best practice all PGDs should be audited on a 6 monthly basis.

Name and post of Designated Lead person within each practice/clinic base:			
Location/Clinic Base:		Date of audit:	
Tick as appropriate. If 'no', state action required	Y	N	Action
Is the PGD or PGDs utilised within the clinical area?			
Has the PGD or PGDs been reviewed within the 2 year limit?			
Do the managers listed on the PGD or PGDs hold a current list of authorised staff?			
Are all staff authorised to work under the PGD or PGDs members of one of the health professions listed in the PGD?			
Do all staff meet the training requirements identified within the PGD?			
Are you confident that all medicines supplied or administered under the PGD or PGDs are stored according to the PGD where this is specified?			
Do the staff working under the PGD or PGDs have a copy of the PGD which has governance sign off and is in date and, available for reference at the time of consultation?			
Where the medicine requires refrigeration. (Delete if not required).			
Is there a designated person responsible for ensuring that the cold chain is maintained?			
Is there a record that the fridge temperature has been monitored to required levels?			
If there is regular and sustained reliance on PGDs for service provision has a Non Medical Prescribing approach been considered as an alternative? (Please note reasons for either a Y/N response).			

Name:	Date of audit:

Keep copies of completed audits alongside your PGD for local reference.

Doxycycline 100mg capsules

Please retain at local level and ensure audit forms are readily available as they may be required for clinical governance audit purposes.